



Hauser-Raspe
Foundation



Quantum Simulation

(...as a tool for computation)

Wolfgang Lechner

QIP 2018 lectures



Hauser-Raspe
Foundation



1. Introduction
2. Adiabatic Theorem
3. Quantum Annealing
4. Entanglement in adiabatic processes
-
5. Parity constraints
6. QRAM
7. Implementations


1. Introduction

Quantum Computing

- Entanglement
 - Superposition
 - Measurement
- ... as a resource.

Universal Quantum Computer

Gate Models



One-Way Quantum Computing

Raussendorf, Briegel PRL (2001).

P. Walther et. al Nature (2005).


Adiabatic Quantum Computing

Fahri et. al., Science (2001).

Quantum Simulation

e.g. Hubbard Model

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$



D Jaksch, et. al. PRL 81, 3108 (1998).

M. Greiner, et. al. Nature, 415 (2002).


Quantum Simulation

Principle: Mimic a quantum system with another controllable quantum system.


Hubbard Models

Theoretical Proposal

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$




D Jaksch, et. al. PRL 81, 3108 (1998).




Non-equilibrium dynamics, Quantum Soft Matter ...

Quantum Glass




WL and P. Zoller, PRL (2013).

Quantum Hexatic




WL, Büchler, P. Zoller, PRL (2014).

Liquid Crystals in Optical Cavities




WL, Habraken, Kiesel, Aspelmeyer, P. Zoller, PRL (2013).

Logarithmic potentials in Rydberg atoms




WL and P. Zoller, PRL (2015).

Supersolid from ultrasoft interactions




Nature Comm. (2015).

Experiment




M. Greiner, et. al. Nature, 415 (2002).

Ultracold Molecules



Jun Ye, Group

Rydberg Atoms




Bloch group Munich
Greiner group Harvard
Saffman group Madison

Quantum Computing

- Entanglement
 - Superposition
 - Measurement
- ... as a resource.

Universal Quantum Computer

Gate Models



One-Way Quantum Computing

Raussendorf, Briegel PRL (2001).

P. Walther et. al Nature (2005).

Adiabatic Quantum Computing

Fahri et. al., Science (2001).

Quantum Simulation

e.g. Hubbard Model

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$



D Jaksch, et. al. PRL 81, 3108 (1998).


M. Greiner, et. al. Nature, 415 (2002).

Universal Quantum Computing - Gate Models



Principle: Universal gates from single qubit unitary and CNOT.

$$|\psi\rangle = U|\psi_0\rangle = U_1U_2U_3\dots|\psi\rangle$$


Single qubit gate



CNOT Gate




Ions




Cirac Zoller PRL 74, 4091 (1995).
Exp: R. Blatt. (2003).

Rydberg Atoms



Saffman group PRL 104, 010503 (2010)

Superconducting Qubits



Mooij group Delft Nature (2007).
J. Martinis, Santa Barbara

Quantum Computing

- Entanglement
 - Superposition
 - Measurement
- ... as a resource.


Universal Quantum Computer

Quantum Simulation

e.g. Hubbard Model

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$

Gate Models




One-Way Quantum Computing

Raussendorf, Briegel PRL (2001).

P. Walther et. al Nature (2005).

Adiabatic Quantum Computing

Fahri et. al., Science (2001).



D Jaksch, et. al. PRL 81, 3108 (1998).

M. Greiner, et. al. Nature, 415 (2002).


Universal Quantum Computing - One Way QC

Principle: Start in a entangled state, get result form consecutive measurements.

Theory:

R .Raussendorf and H. Briegel, PRL 86, 5188 (2001).

Experiment:



P. Walther, Resch, Rudolph, Schenck, Weinfurter, V. Vedral, M. Aspelmeyer and A. Zeilinger Nature.434, 176 (2005).

Quantum Computing

- Entanglement
 - Superposition
 - Measurement
- ... as a resource.

Universal Quantum Computer

Quantum Simulation

e.g. Hubbard Model

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$

Gate Models



One-Way Quantum Computing

Raussendorf, Briegel PRL (2001).

P. Walther et. al Nature (2005).

Adiabatic Quantum Computing

Fahri et. al., Science (2001).



D Jaksch, et. al. PRL 81, 3108 (1998).

M. Greiner, et. al. Nature, 415 (2002).


Quantum Computing

- Entanglement
 - Superposition
 - Measurement
- ... as a resource.

Universal Quantum Computer

Quantum Simulation

Gate Models



One-Way
Quantum
Computing

Raussendorf, Briegel PRL (2001).


P. Walther et. al Nature (2005).

Adiabatic Quantum
Computing

Farhi et. al., Science (2001).

e.g. Hubbard Model

$$H = -t \sum_{\langle i,j \rangle} a_i^\dagger a_j + \frac{U}{2} \sum_i \hat{n}_i (\hat{n}_i - 1) - \mu \sum_i \hat{n}_i$$



D Jaksch, et. al. PRL 81, 3108 (1998).

M. Greiner, et. al. Nature, 415 (2002).

2. The Adiabatic Theorem

Adiabatic Theorem

A physical system **remains** in its **instantaneous eigenstate** if a given perturbation is acting on it **slowly enough** and if **there is a gap** between the eigenvalue and the rest of the Hamiltonian's spectrum.

M. Born and V. A. Fock "Beweis des Adiabatenatzes" Zeitschrift für Physik A. **51** 165 (1928).

Avoided Level Crossings

Two-level system


$$H = \begin{pmatrix} E_1 & 0 \\ 0 & E_2 \end{pmatrix}$$

Eigenvalues
 E_1 and E_2

Eigenvectors
 $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ and $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$

Time-dependent

$$H(t) = \begin{pmatrix} E_1(t) & 0 \\ 0 & E_2(t) \end{pmatrix}$$




Avoided Level Crossings

Two-level system with coupling

$$H(t) = \begin{pmatrix} E_1 & W \\ W^* & E_2 \end{pmatrix}$$

e.g. Rabi Problem

$$\hbar \begin{pmatrix} -\Delta & \frac{\Omega}{2} e^{-i\varphi} \\ \frac{\Omega}{2} e^{-i\varphi} & 0 \end{pmatrix}$$



Eigenvalues

$$E_+ = \frac{1}{2}(E_1 + E_2) + \frac{1}{2}\sqrt{(E_1 - E_2)^2 + 4|W|^2}$$

$$E_- = \frac{1}{2}(E_1 + E_2) - \frac{1}{2}\sqrt{(E_1 - E_2)^2 + 4|W|^2}$$

Eigenstates (“dressed states”)

$$|+\rangle = \frac{1}{\sqrt{2}}(e^{i\phi}|0\rangle + |1\rangle)$$


$$\text{with } \tan \phi = -\frac{\Omega}{\Delta}$$

$$|-\rangle = \frac{1}{\sqrt{2}}(-e^{i\phi}|0\rangle + |1\rangle)$$

Avoided Level Crossings

Two-level system with coupling

$$H(t) = \begin{pmatrix} E_1 & W \\ W^* & E_2 \end{pmatrix}$$



“avoided crossing”

Adiabatic Theorem

Time-dependent Schrödinger Equation

$$i\hbar \frac{\partial}{\partial t} \Psi(x, t) = H \Psi(x, t)$$

General Solution as a function of instantaneous eigenstates

$$\Psi(x, t) = \sum_n c_n \Psi_n(x, t) = \sum_n c_n \psi_n(x) e^{-iE_n t/\hbar}$$

with $\Psi_n(x, t) = \psi_n(x) e^{-iE_n t/\hbar}$

For each instantaneous time slice we have

$$H(t)\psi_n(x, t) = E_n(t)\psi_n(x, t) \text{ with } \langle \psi_n(t) | \psi_m(t') \rangle = \delta_{nm} \delta(t - t')$$

time-independent Schrödinger equation with time as parameter

Adiabatic Theorem

Time-dependent Schrödinger Equation

$$i\hbar \frac{\partial}{\partial t} \Psi(x, t) = H \Psi(x, t)$$

with “dynamical Phase factor”

$$\Psi(t) = \sum_n c_n(t) \psi_n(t) e^{i\theta_n(t)} \quad \theta_n(t) = -\frac{1}{\hbar} \int_0^t E_n(t') dt'$$

Plug this into Schrödinger equation

$$i\hbar \sum_n (\dot{c}_n \psi_n + c_n \dot{\psi}_n + \cancel{i c_n \psi_n \dot{\theta}_n}) e^{i\theta_n} = \sum_n \cancel{c_n H \psi_n} e^{i\theta_n}$$
$$\dot{\theta}_n = -E_n / \hbar$$

$$\Rightarrow \sum_n \dot{c}_n \psi_n e^{i\theta_n} = - \sum_n c_n \dot{\psi}_n e^{i\theta_n}$$

Adiabatic Theorem

$$\sum_n \dot{c}_n \psi_n e^{i\theta_n} = - \sum_n c_n \dot{\psi}_n e^{i\theta_n}$$

multiply with $\langle \psi_m |$ from both sides

$$\dot{c}_m(t) = - \sum_n c_n \langle \psi_m | \dot{\psi}_n \rangle e^{i(\theta_n - \theta_m)} \quad (\text{with } \langle \psi_m | \psi_n \rangle = \delta_{nm})$$

evaluating $\dot{\psi}_n$ from the Schrödinger equation

$$\dot{c}_m(t) = -c_m \langle \psi_m | \dot{\psi}_m \rangle - \sum_{n \neq m} c_n \frac{\langle \psi_m | \dot{H} | \psi_n \rangle}{E_n - E_m} e^{i(\theta_n - \theta_m)}$$

↑
This is the adiabatic approximation

$$\text{Consequence: } \dot{c}_m(t) = -c_m \langle \psi_m | \dot{\psi}_m \rangle$$

Adiabatic Theorem

As a consequence of the approximation, matrix elements from m to n vanish.

$$\dot{c}_m(t) = -c_m \langle \psi_m | \dot{\psi}_m \rangle$$

with the formal solution:

$$c_m(t) = c_m(0) \exp\left[-\int_0^t \langle \psi_m(t') | \dot{\psi}_m(t') \rangle dt'\right] = c_m(0) e^{i\gamma_m(t)}$$

where $\gamma_m(t)$ is the geometric phase factor

$$\gamma_m(t) = i \int_0^t \langle \psi_m(t') | \dot{\psi}_m(t') \rangle dt'$$

plugging $c_m(t)$ into the solution of the Schrödinger equation

$$\Psi_n(t) = c_n(t) \psi_n(t) e^{i\gamma_n(t)} = \psi_n(t) e^{i\theta_n(t)} e^{i\gamma_n(t)}$$

Berry Phase


Landau Zener

C. Zener, Proc. Royal Soc. A 137 696(1923). A Vutar, arXiv:1001.3322 (2010).

$$\mathbf{H} = \begin{pmatrix} 0 & \Omega^\dagger e^{i\omega t} \\ \Omega e^{-i\omega t} & \omega_0 \end{pmatrix}$$

$$\Delta = \omega - \omega_0$$

$$\dot{\Delta} = \text{const}$$




Transition Rate

$$\Gamma = \Omega^2 \frac{\gamma}{\Delta^2 + \gamma^2/4}$$

$$\gamma \sim \sqrt{\frac{\dot{\Delta}}{4\pi}}$$

Landau Zener

$$P_0(t + \delta t) = [1 - \Gamma(t)\delta t]P_0(t) \approx e^{-\Gamma(t)\delta t}P_0(t)$$



Integrating from initial to final time


$$\begin{aligned} P(t_f) &= \exp\left[-\int_{t_i}^{t_f} \Gamma(t)dt\right] \\ &= \exp\left\{-\frac{2\Omega^2}{\dot{\Delta}}\left[\text{atan}\left(\frac{\Delta(t_f)}{\gamma/2}\right) - \text{atan}\left(\frac{\Delta(t_i)}{\gamma/2}\right)\right]\right\} \end{aligned}$$

At the level crossing $\Delta \rightarrow 0$


$$P(t_f) = e^{-\frac{2\pi\Omega^2}{\dot{\Delta}}}$$

Landau Zener Formula

Landau Zener



Picture taken from J. R Johnson et. al. QuTip Package



Landau Zener

Dimensionless schedule s from $0 \rightarrow 1$

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = H(t) |\psi(t)\rangle \Rightarrow \frac{d}{ds} |\psi(s)\rangle = -i\tau(s) H(s) |\psi(s)\rangle$$

introducing the “delay factor” $\tau(s)$


$$\tau(s) \gg \frac{|\frac{d}{ds} H(s)|}{g(s)^2} \quad \text{Landau Zener condition with gap } g(s)$$

Total time of the calculation

$$T = \int_{s=0}^1 \tau(s) ds$$

Landau Zener

Many-Body Landau Zener



Approximation:
sequence of LZ transitions

$$P_{LZ} = e^{-\frac{|G|^2}{v}}$$

Gap is estimated from off-diagonal matrix elements:

$$|G| \sim |\langle n|H|m\rangle|$$

3. Quantum Annealing

Adiabatic Quantum Computing

Nishimori Hidetoshi Phys. Rev. B (1998).

E. Farhi et. al., Science 292, 472 (2001).

$$H(t) = \left(1 - \frac{t}{T}\right) H(0) + \frac{t}{T} H_p$$

Trivial initial Hamiltonian


Problem Hamiltonian

1. Prepare System in $|\psi_0\rangle$
which is the ground state of $H(0)$

2. Slowly evolve the system with

$$|\psi(t)\rangle = e^{-i \int_0^t H(t) dt} |\psi_0\rangle$$

3. Due to the adiabatic theorem, the final state is the ground state of H_p



$$H_0 = \sum_i^N \sigma_x^{(i)}$$

$$H_p = \sum_{i < j} J_{ij} \sigma_z^{(i)} \sigma_z^{(j)}$$

Adiabatic Quantum Computing

Nishimori Hidetoshi Phys. Rev. B (1998).

E. Farhi et. al., Science 292, 472 (2001).

$$H(t) = \left(1 - \frac{t}{T}\right) H(0) + \frac{t}{T} H_p$$

Trivial initial Hamiltonian


Problem Hamiltonian

1. Prepare System in $|\psi_0\rangle$
which is the ground state of $H(0)$

2. Slowly evolve the system with

$$|\psi(t)\rangle = e^{-i \int_0^t H(t) dt} |\psi_0\rangle$$

3. Due to the adiabatic theorem, the final state is the ground state of H_p



$$H_0 = \sum_i^N \sigma_x^{(i)}$$

$$H_p = \sum_{i < j} J_{ij} \sigma_z^{(i)} \sigma_z^{(j)}$$

Adiabatic Quantum Computing

Nishimori Hidetoshi Phys. Rev. B (1998).

E. Farhi et. al., Science 292, 472 (2001).

$$H(t) = \left(1 - \frac{t}{T}\right) H(0) + \frac{t}{T} H_p$$

Trivial initial Hamiltonian


Problem Hamiltonian

1. Prepare System in $|\psi_0\rangle$
which is the ground state of $H(0)$

2. Slowly evolve the system with

$$|\psi(t)\rangle = e^{-i \int_0^t H(t) dt} |\psi_0\rangle$$

3. Due to the adiabatic theorem, the final state is the ground state of H_p



$$H_0 = \sum_i^N \sigma_x^{(i)}$$

$$H_p = \sum_{i < j} J_{ij} \sigma_z^{(i)} \sigma_z^{(j)}$$

Example: Quantum Adiabatic Search

J. Roland and N. J. Cerf Phys. Rev. A 65, 042308 (2001).

W. Dam, et. al. arxiv:quant-ph/0206003 (2008).

$$f : \{0, 1\}^n \rightarrow \mathbb{R} \qquad f(u) = \begin{cases} 0, & \text{if } u \in \{0, 1\}^n. \\ 1, & \text{otherwise.} \end{cases}$$

where u is the search string.

where u is the search string.

Hamiltonian for quantum annealing:

Spectrum

$$H_u = \sum_{z \in \{0, 1\}^n \setminus u} |z\rangle\langle z|$$

1	————	all other strings
0	————	search string

Example: Quantum Adiabatic Search

W. Dam, et. al. arxiv:quant-ph/0206003 (2008).

initial Hamiltonian

$$H_0 = \sum_{z \in \{0,1\}^n \setminus \{0\}^n} |z\rangle\langle z|$$

final Hamiltonian


$$H_u = \sum_{z \in \{0,1\}^n \setminus u} |z\rangle\langle z|$$

Adiabatic protocol

$$H(t) = \left(1 - \frac{t}{T}\right) H_0 + \frac{t}{T} H_u$$

Gap as a function of $s = \frac{t}{T}$

$$g(s) = \sqrt{\frac{2^n + 4(2^n - 1)(s^2 - s)}{2^n}}$$



Example: Quantum Adiabatic Search

W. Dam, et. al. arxiv:quant-ph/0206003 (2008).

$$\text{Minimum at } s^* = \frac{t}{T} = \frac{1}{2} \quad g(s^*) = \sqrt{\frac{2^n + 4(2^n - 1)(-\frac{1}{4})}{2^n}} = \frac{1}{\sqrt{2^n}}$$

Naive :

$$T \approx \mathcal{O}(g(s^*)^{-2}) \sim \mathcal{O}(2^n) = \mathcal{O}(N)$$

Inverse gap schedule


$$\begin{aligned} T &= \int_{s=0}^1 \frac{ds}{g(s)^2} = \int_{s=0}^1 \frac{2^n}{2^n + (2^n - 1)(s^2 - s)} ds \\ &= \frac{2^n \arctan(\sqrt{2^n - 1})}{\sqrt{2^n - 1}} \sim \mathcal{O}(\sqrt{2^n}) = \mathcal{O}(\sqrt{N}) \end{aligned}$$

Quadratic Grover Speedup

further reading: D. Wild et.al. Phys. Rev. Lett. **117**, 150501 (2016).

Example: Quantum Optimization

Graph Partitioning




$$G = (V, E)$$

We ask: What is a partition of the set V into two subsets of equal size $N/2$ such that the number of edges connecting the two subsets is minimized?

Example: Quantum Optimization

Graph Partitioning




$$G = (V, E)$$

We ask: What is a partition of the set V into two subsets of equal size $N/2$ such that the number of edges connecting the two subsets is minimized?

Example: Quantum Optimization

Graph Partitioning



$$G = (V, E)$$

We ask: What is a partition of the set V into two subsets of equal size $N/2$ such that the number of edges connecting the two subsets is minimized?

Strategy: Cost-function = Energy

$$\sigma_i = +1 \quad \text{○}$$

$$\sigma_i = -1 \quad \text{○}$$

$$H_p = C_A \left(\sum_{n=1}^N \sigma_i \right)^2 + C_B \sum_{\{i,j\} \in E} \frac{1 - \sigma_i \sigma_j}{2}$$

Two subsets have the same size.

Connection between two sets is minimal.

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$

Quantum Optimization

NP-complete problems = Infinite range spin glass

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$

List of NP-complete problems:

A. Lucas, *Frontiers in Physics* **2**, 00005 (2014).

- Traveling Salesman
- Knapsack
- 3SAT
- Number Partitioning
- Graph Coloring
- Minimal Spanning Tree
- Graph Partition
- ... and many more

Quantum Optimization

NP-complete problems = Infinite range spin glass

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$

Scientific optimization problems:


Protein folding



e.g. beta-lactoglobulin (milk protein)
Picture: Peter Bolhuis

I. Coluzza, et.al. Biophys. J. (2007).

Quantum Chemistry




Picture: E. Meijer, University of Amsterdam.

Adiabatic Quantum computing: A. Perdomo-Ortiz et. al.,
Sci. Rep. 2, 571 (2012).

Adiabatic Quantum Computing: R. Babbush et. al.,
Sci. Rep. 4, 6603 (2014).

4. Entanglement in adiabatic processes



Entanglement in an Adiabatic Protocol



Entanglement in an Adiabatic Protocol

P. Hauke, L. Bonnes, M. Heyl and W. Lechner, Front. Phys. 3, 21 (2015).

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$




Entanglement?

Product States

Entanglement measures

Van Neumann Entropy

Relative entropy of two subsystems A and B



Entanglement Measure

$$S_A = -\text{Tr} [\rho_A \log(\rho_A)]$$

$$\rho_A = \text{Tr}_B \rho$$

Take Maximum of all combinations of A and B and maximum at each time $S_2^{\max} = \max_t [S_2^{\max}(t)]$

Features


- + well-established, simple to calculate
- only closed systems
- only block-pair entanglement

Entanglement measures

Quantum Fisher Information

P. Hyllus et.al., Phys. Rev. A 85, 022321 (2012).

Phase estimation experiment:

$$\rho(\theta) = e^{-i\theta H} \rho e^{i\theta H}$$


Want to measure unknown phase

Estimate θ from m independent POVM (positive operator values measurements) $\{E_\mu\}$

$\theta_{est}(\{\mu_i\}_m)$ with $\{\mu_i\}_m = \{\mu_1, \mu_2, \dots, \mu_m\}$
m measurement outcomes μ

Average

$$\langle \theta_{est} \rangle$$

Standard deviation

$$(\Delta \theta_{est})^2 = \langle \theta_{est}^2 \rangle - \langle \theta_{est} \rangle^2$$

Entanglement measures

P. Hyllus et.al., Phys. Rev. A 85, 022321 (2012).

Standard Deviation is limited by the bounds

$$\Delta\theta_{est} \geq \frac{1}{\sqrt{mF_Q}}$$

Quantum Fisher Information

$$F_Q[\rho, H] = 2 \sum_{l,l'} \frac{(\lambda_l - \lambda_{l'})^2}{\lambda_l + \lambda_{l'}} |\langle l|H|l'\rangle|$$

for any mixed state

$$\rho = \sum_l \lambda_l |l\rangle\langle l| \quad \text{with} \quad \sum_l \lambda_l = 1 \quad \lambda_l > 0$$

Entanglement measures

P. Hyllus et.al., Phys. Rev. A 85, 022321 (2012).

- A pure state is **k-producible** if it can be written as

$$|\psi_{\text{k-prod}}\rangle = \bigotimes_{l=1}^M |\psi_l\rangle \text{ where } |\psi_l\rangle \text{ is a state of } N_l \leq k \text{ particles} \\ \text{with } \sum_l^M N_l = N$$

- A pure state is **k-particle entangled** if it is k-producible but not (k-1)-producible.

- A mixed state is k-producible if it can be written as

$$\rho_{\text{k-prod}} = \sum_l p_l |\psi_{\text{k-prod}}\rangle \langle \psi_{\text{k-prod}}|$$

Entanglement measures

P. Hyllus et.al., Phys. Rev. A 85, 022321 (2012).

Quantum Fisher Information bounds k-particle entanglement

$$F_Q \leq \lfloor \frac{N}{k} \rfloor k^2 + \left(N - \lfloor \frac{N}{k} \rfloor k \right)^2$$

where $\lfloor \cdot \rfloor$ denotes a floor operation

Features


- + measure for multipartite entanglement
- + also density matrices
- more involved to measure

Entanglement and Success Probability

P. Hauke, L. Bonnes, M. Heyl and W. Lechner, Front. Phys. 3, 21 (2015).

Instantaneous probability to be in the ground state

$$P(t) = |\langle \psi(t) | \psi_{gs} \rangle|^2.$$



Success Probability is the ground state overlap at the final time

$$P_{\text{succ}} = |\langle \psi(T) | \psi_{gs} \rangle|^2.$$

Entanglement and Success Probability

P. Hauke, L. Bonnes, M. Heyl and W. Lechner, Front. Phys. 3, 21 (2015).

A. Polkovnikov, Annals of Physics 326, 486 (2011).

Ideal solution

$$|\psi(T)\rangle = |s_A^* s_B^*\rangle$$

Finite speed

$$\rho_A = \text{Tr}_B \rho$$

Entropy from population of “wrong” states

$$S_A \leq S_A^d = -\text{tr}[\rho_A^d \log(\rho_A^d)] = -\sum_{\mu=1}^{d_A} p_\mu \log(p_\mu)$$

ρ_A^d only diagonal elements of ρ_A

$d_A = 2^l$ number of spins in A


p_μ Eigenvalues of ρ_A^d

Entanglement Entropy as a function of the Success probability

$$S_l^A \leq -P \log(P) - (1 - P) \log(1 - P) + (1 - P) \log(2^l - 1)$$

Entanglement and Success Probability



P. Hauke, L. Bonnes, M. Heyl and W. Lechner, Front. Phys. 3, 21 (2015).



Entanglement and Success Probability

P. Hauke, L. Bonnes, M. Heyl and W. Lechner, Front. Phys. 3, 21 (2015).

Individual Run: Connection between irreversibility and entanglement



5. Quantum Annealing with Parity Constraints

WL, P. Hauke, P. Zoller, Science Advances 1, 1500838 (2015).


Adiabatic Quantum Annealing

Spin Glass paradigm of AQC

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$

Superconducting Qubits

Chimera-graphs




(D-Wave Systems)

M. W. Johnson et. al. Supercond. Sci. Technol. **23** 065004 (2010).

Z. Bian, et. al. Phys. Rev. Lett. **111**, 130505 (2013).


New platforms for Quantum Annealing

Ultracold atoms in optical lattices




Bloch, Munich

Rydberg atoms




Saffman, Madison

Hybrid Ion-traps



S. Benjamin, Oxford


Transmons



Filipp, IBM


A. Wallraff

Ions in surface traps



Blatt, Innsbruck


Polar Molecules



J. Ye, Boulder

Parity Based Quantum Annealing


Spin glass paradigm



$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$



Parity paradigm




$$\mathcal{H}(t) = A(t) \sum_{i=1}^K b_i \sigma_x^{(i)} + B(t) \sum_{i=1}^K J_i \sigma_z^{(i)} + C(t) \sum_{l=1}^{K-N} C_l$$

WL, P. Hauke, P. Zoller, Science Advances 1, 1500838 (2015).

- encode **all-to-all interaction matrix** in local fields
- architecture comprises **only local interaction** on 2D geometry
- can be realized in current **qubit platform**
(e.g. flux qubits, transmon qubits, ultracold atoms, hybrids systems,...)
- **intrinsic error tolerance** from redundant encoding


All-to-all programmable architecture

Transformation to parity variables.




Physical qubits $\hat{\sigma}_z^{(k)}$ represent parity of logical qubits.

All-to-all programmable architecture



$$\mathcal{H} = \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} = \sum_{i=1}^K J_i \hat{\sigma}_z^{(i)}$$

All-to-all programmable architecture




$$\mathcal{H} = \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} = \sum_{i=1}^K J_i \hat{\sigma}_z^{(i)}$$

N degrees of freedom

$$K = \frac{N(N-1)}{2}$$

degrees of freedom


All-to-all programmable architecture




$$\mathcal{H} = \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} = \sum_{i=1}^K J_i \hat{\sigma}_z^{(i)} + \sum_{l=1}^{K-N} C_l$$

K-N constraints

All-to-all programmable architecture



$$\mathcal{H} = \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} = \sum_{i=1}^K J_i \hat{\sigma}_z^{(i)} + \sum_{l=1}^{K-N} C_l$$




K-N constraints

What are these constraints?

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1			

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1		

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	1

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	1
0	0	0	1	1	1

Constraints


Conditions on closed loops



$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	1
0	0	0	1	1	1
0	0	1	1	0	0

Constraints


Conditions on closed loops




$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$	$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	1
0	0	0	1	1	1
0	0	1	1	0	0
...			...		
			1	1	0

Constraints

Conditions on closed loops





$\sigma_z^{(1)}$	$\sigma_z^{(2)}$	$\sigma_z^{(3)}$			
			$\hat{\sigma}_z^{(12)}$	$\hat{\sigma}_z^{(13)}$	$\hat{\sigma}_z^{(23)}$
1	1	1	1	1	1
0	0	0	1	1	1
0	0	1	1	0	0
...			...		
			1	1	0

In **each closed loop**, the number of **spin-down** has to be an **even** number or 0.



Constraints

Logical Qubits

Physical Qubits




$$C_l = -C \hat{\sigma}_z^{(12)} \hat{\sigma}_z^{(23)} \hat{\sigma}_z^{(13)}$$




$$C_l = -C \hat{\sigma}_z^{(12)} \hat{\sigma}_z^{(23)} \hat{\sigma}_z^{(34)} \hat{\sigma}_z^{(14)}$$

Constraints



Constraints



$s=5$

$s=4$

$s=3$


$s=2$

$s=1$



23

Constraints



$s=5$

$s=4$

$s=3$

$s=2$

$s=1$





13





23


Constraints




Constraints



Constraints





Number of spin-down
is even





$$C_l = -C \left(\sum_{m=n,e,s,w} \tilde{\sigma}_z^{(l,m)} + S_z^l \right)^2 = -C \tilde{\sigma}_z^{(l,n)} \tilde{\sigma}_z^{(l,e)} \tilde{\sigma}_z^{(l,s)} \tilde{\sigma}_z^{(l,w)}$$



Constraints




Constraints





Constraints



Constraints



Constraints




Parity AQC

$$\mathcal{H}(t) = A(t) \sum_{i=1}^K b_i \sigma_x^{(i)} + B(t) \sum_{i=1}^K J_i \sigma_z^{(i)} + C(t) \sum_{l=1}^{K-N} C_l$$



$$K = \frac{N(N-1)}{2}$$

$$C_l = -C \tilde{\sigma}_z^{(l,n)} \tilde{\sigma}_z^{(l,e)} \tilde{\sigma}_z^{(l,s)} \tilde{\sigma}_z^{(l,w)}$$



Spectrum



$$H(t) = A(t) \sum_{k=1}^K h_k \tilde{\sigma}_x^{(k)} + B(t) \left[\sum_{k=1}^K J_k \tilde{\sigma}_z^{(k)} + \sum_{l=1}^{K-N} C_l \right]$$



Magnetic Field term

$$H = \sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)}$$


Add 1 “row” of qubits.




Three-Body interactions



$$H = \sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} + \sum_{i=1}^N \sum_{j=1}^i \sum_{k=1}^j J_{ijk} \sigma_z^{(i)} \sigma_z^{(j)} \sigma_z^{(k)}$$

i	j	k	
↑	↑	↑	= 1
↑	↑	↓	= 0
↑	↓	↑	= 0
↑	↓	↓	= 1
↓	↑	↑	= 0
↓	↑	↓	= 1
↓	↓	↑	= 1
↓	↓	↓	= 0


 #0 = 0, 2 or 4




Readout





Readout






Determining and Non-determining Readouts






Determining and Non-determining Readouts







Determining and Non-determining Readouts







Determining and Non-determining Readouts



Determining and Non-determining Readouts




Determining and Non-determining Readouts




Error correction

Spin glass annealing



Gauge field annealing



Number of qubits

$$N$$

$$\frac{N(N-1)}{2}$$

Individual Qubit error rate

$$\Gamma$$

$$\Gamma$$


Success Probability

$$P_s = 1 = \sum_i^N \Gamma \tau I_i = 1 - \Gamma \tau I_s N$$


$$P_s = 1 - \Gamma \tau I_g \frac{N(N-1)}{2}$$

Error correction

Spin glass annealing



Gauge field annealing



Success Probability


$$P_s = 1 - \Gamma \tau N I_s$$

$$I_s = 1$$


$$P_s = 1 - \Gamma\tau \frac{N(N-1)}{2} I_g$$

Error correction

Spin glass annealing



Gauge field annealing




Success
Probability

$$P_s = 1 - \Gamma\tau NI_s$$

$$I_s = 1$$

$$P_s = 1 - \Gamma\tau \frac{N(N-1)}{2} I_g$$



— $\Gamma\tau \frac{N(N-1)}{2}$

— I_g

--- $\Gamma\tau \frac{N(N-1)}{2} I_g$

$$P_s = 1 - \Gamma\tau(N-1)$$

Error correction

Fernando Pastawski, John Preskill, arXiv/1511.00004 (2015).

Belief propagation algorithm

Parity is preserved for any closed loop: $0 = (12) \oplus (23) \oplus (13) = (12) \oplus (24) \oplus (14) = \dots = (12) \oplus (2N) \oplus (1N)$



Estimate value of (12) from

$$g_{12} = (23) \oplus (13) = (24) \oplus (14) = \dots \quad \text{loops of 3}$$

$$g_{12} = (23) \oplus (34) \oplus (14) = (23) \oplus (35) \oplus (15) = \dots = \text{loops of 4}$$

$$g_{12} = (23) \oplus (34) \oplus (45) \oplus (15) = \dots \quad \text{loops of 5}$$

■ ■ ■




6. QRAM

Many-Body Spin Superposition

List of Bit-Strings

1011
1100
1111
...



$$|\psi\rangle = a_1|1011\rangle + a_2|1100\rangle + a_3|1111\rangle$$


Quantum Superposition

Many-Body Spin Superposition

List of Bit-Strings

Quantum Superposition

1011
1100
1111
...



$$|\psi\rangle = a_1|1011\rangle + a_2|1100\rangle + a_3|1111\rangle$$



This Talk!


arXiv:1708.02533 (2017).

Adiabatic Theorem

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$


H. Nishimori (1998).

E. Farhi (2001).




Adiabatic Theorem

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$




Adiabatic Theorem

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$



Fair Sampling

$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$



$$|\psi\rangle = a_1 | \text{state (1)} \rangle + a_2 | \text{state (2)} \rangle + a_3 | \text{state (3)} \rangle$$


Exponential bias in amplitudes

H. Katzgraber et.al. Phys. Rev. Lett. **118**, 070502 (2017).

Can we make these amplitudes tunable?


Parity Constraints

Spin glass paradigm



$$\mathcal{H}(t) = A(t) \sum_i^N b_i \sigma_x^{(i)} + B(t) \left[\sum_{i=1}^N h_i \sigma_z^{(i)} + \sum_{i=1}^N \sum_{j=1}^i J_{ij} \sigma_z^{(i)} \sigma_z^{(j)} \right]$$

Parity constraints





$$\mathcal{H}(t) = A(t) \sum_{i=1}^K b_i \sigma_x^{(i)} + B(t) \sum_{i=1}^K J_i \sigma_z^{(i)} + C(t) \sum_{l=1}^{K-N} C_l$$

- comprises **problem-independent, local interaction** -> **Gauge invariance**

Spectrum

$$H(t) = A(t) \sum_{i=1}^K \sigma_i^x + B(t) \sum_{i=1}^K J_i \sigma_i^z + C(t) \sum_{p=1}^{K-N+1} C_p \sigma_{n_p}^z \sigma_{w_p}^z \sigma_{s_p}^z \sigma_{e_p}^z.$$




Gauge Invariant

Invariant
under
variation of
individual C_p

Spectrum

$$H(t) = A(t) \sum_{i=1}^K \sigma_i^x + B(t) \sum_{i=1}^K J_i \sigma_i^z + C(t) \sum_{p=1}^{K-N+1} C_p \sigma_{n_p}^z \sigma_{w_p}^z \sigma_{s_p}^z \sigma_{e_p}^z.$$




Strategy: make use of gauge degrees of freedom to tune the quantum phase of the paths.

Effective Theory

$$H(t) = A(t) \sum_{i=1}^K \sigma_i^x + B(t) \sum_{i=1}^K J_i \sigma_i^z + C(t) \sum_{p=1}^{K-N+1} C_p \sigma_{n_p}^z \sigma_{w_p}^z \sigma_{s_p}^z \sigma_{e_p}^z.$$

Adiabatic process



Adiabatic manifold:

- Projection

$$\mathcal{P} = \sum_{n=1}^M |z_n\rangle\langle z_n|$$

$$Q = 1 - \mathcal{P}$$


- Schrieffer-Wolff

- Perturbation Theory

Effective Theory

$$H(t) = A(t) \sum_{i=1}^K \sigma_i^x + B(t) \sum_{i=1}^K J_i \sigma_i^z + C(t) \sum_{p=1}^{K-N+1} C_p \sigma_{n_p}^z \sigma_{w_p}^z \sigma_{s_p}^z \sigma_{e_p}^z.$$

Adiabatic process



$$H_{\text{eff}}(t) = \begin{pmatrix} e_1 t^{-2} & g_{12} t^{-h_{12}} & g_{13} t^{-h_{13}} \\ g_{21} t^{-h_{21}} & e_2 t^{-2} & g_{23} t^{-h_{23}} \\ g_{31} t^{-h_{31}} & g_{32} t^{-h_{32}} & e_3 t^{-2} \end{pmatrix}$$


Matrix Elements

Spin-Model

1100


1111

Lattice Gauge



Count Feynman Paths!


Feynman Paths



Matrix Elements

$$g_{12} = \frac{1}{2^3} \left(\begin{aligned} & - \frac{\frac{1}{C_2+C_3} + \frac{1}{1+C_1+C_3}}{1+C_1+C_2} + \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{1+C_1+C_3}}{C_2} - \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{C_2+C_3}}{1-C_1} \\ & \frac{C_3}{C_3} \\ & - \frac{-\frac{1}{C_2+C_3} - \frac{1}{C_3}}{C_2} - \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{C_2+C_3}}{1-C_1} - \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{C_3}}{1-C_1-C_2} \\ & \frac{1-C_1-C_3}{1-C_1-C_3} \\ & + \frac{-\frac{1}{1+C_1+C_3} + \frac{1}{C_3}}{1+C_1} + \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{1+C_1+C_3}}{C_2} - \frac{\frac{1}{1-C_1-C_2-C_3} - \frac{1}{C_3}}{1-C_1-C_2} \\ & \frac{C_2+C_3}{C_2+C_3} \\ & + \frac{-\frac{1}{1+C_1+C_3} + \frac{1}{C_3}}{1+C_1} - \frac{\frac{1}{C_2+C_3} + \frac{1}{C_3}}{C_2} - \frac{\frac{1}{C_2+C_3} + \frac{1}{1+C_1+C_3}}{1+C_1+C_2} \\ & \frac{1+C_1+C_2+C_3}{1+C_1+C_2+C_3} \end{aligned} \right)$$


Example




Fair Sampling

$$|\psi\rangle = \sqrt{\frac{1}{3}}|1011\rangle + \sqrt{\frac{1}{3}}|1100\rangle + \sqrt{\frac{1}{3}}|1111\rangle$$

Result: $C_1 = 9.4$ $C_2 = 0.4$ $C_3 = 9.2$




Example






Programmable Weights

$$|\psi\rangle = \sqrt{0.2}|1011\rangle + \sqrt{0.3}|1100\rangle + \sqrt{0.5}|1111\rangle$$

Result: $C_1 = 5.6$ $C_2 = 3.0$ $C_3 = 6.2$



Ergodicity




7.Implementations

New platforms for Quantum Annealing


A. Glätzle, R. van Bijnen, P. Zoller and WL, arXiv:1611.02594 (2016).

Ultracold atoms in optical lattices




Bloch, Munich

Rydberg atoms




Saffman, Madison

Hybrid Ion-traps



S. Benjamin, Oxford


Superconducting Qubits



A. Wallraff


Filipp, IBM

Ions in surface traps



Blatt, Innsbruck

Polar Molecules





J. Ye, Boulder

Martin Leib, P. Zoller, and WL, arXiv:1604.02359(2016).


Strategies for constraint design with Rydberg atoms

4-body constraints




$$H_c = -C\sigma_z^1\sigma_z^2\sigma_z^3\sigma_z^4$$

Unphysical states do not matter



Strategies for constraint design with Rydberg atoms

Andrea Rocchetto, Simon C. Benjamin and Ying Li, Science Advances 2, e1601246 (2016).




$$\sum_i^4 \sigma_z^{(i)}$$

↑

-4, 0, 4

Strategies for constraint design with Rydberg atoms


Andrea Rocchetto, Simon C. Benjamin and Ying Li, Science Advances 2, e1601246 (2016).




$$S = \left(\underset{\substack{\uparrow \\ \text{qutrit}}}{4S_z} + \underset{\substack{\uparrow \\ -4,0,4}}{\sum_i^4 \sigma_z^{(i)}} \right)^2$$

Strategies for constraint design with Rydberg atoms

Andrea Rocchetto, Simon C. Benjamin and Ying Li, Science Advances 2, e1601246 (2016).




$$S = \left(\underset{\substack{\uparrow \\ \text{qutrit}}}{4S_z} + \underset{\substack{\uparrow \\ -4,0,4}}{\sum_i^4 \sigma_z^{(i)}} \right)^2$$




$$S = \left(\underset{\substack{\uparrow \\ \text{qubit}}}{2\sigma_z^{(a)}} + \underset{\substack{\uparrow \\ -2,2}}{\sum_i^4 \sigma_z^{(i)}} \right)^2$$

Odd parity representation




Odd parity representation


Andrea Rocchetto, Simon C. Benjamin and Ying Li, Science Advances 2, e1601246 (2016).




Odd parity representation



$$S = \left(2\sigma_z^{(a)} + \sum_i^4 \sigma_z^{(i)} \right)^2$$











Odd parity representation





$$S = \left(2\sigma_z^{(a)} + \sum_i^4 \sigma_z^{(i)} \right)^2$$

$$2\sigma_z^{(a)}\sigma_z^{(1)} + 2\sigma_z^{(a)}\sigma_z^{(2)} + 2\sigma_z^{(a)}\sigma_z^{(3)} + 2\sigma_z^{(a)}\sigma_z^{(4)} + \sigma_z^{(1)}\sigma_z^{(2)} + \sigma_z^{(1)}\sigma_z^{(3)} + \sigma_z^{(1)}\sigma_z^{(4)} + \sigma_z^{(2)}\sigma_z^{(3)} + \sigma_z^{(2)}\sigma_z^{(4)} + \sigma_z^{(3)}\sigma_z^{(4)}$$




Error Robustness





Neutral Atoms in Optical Lattices

A. Glätzle, R. van Bijnen, P. Zoller and WL, arXiv:1611.02594 (2016).



I. I. Beterov and M. Saffman,
Phys. Rev. A 92, 042710 (2015).




$$\hat{H} = \hat{H}_A^{(1)} \otimes \hat{I} + \hat{I} \otimes \hat{H}_A^{(2)} + \hat{V}_{dd}$$


R. M. W. van Bijnen and T. Pohl
Phys. Rev. Lett. **114**, 243002 (2015)

Neutral Atoms in Optical Lattices

A. Glätzle, R. van Bijnen, P. Zoller and WL, arXiv:1611.02594 (2016).





I. I. Beterov and M. Saffman,
Phys. Rev. A 92, 042710 (2015).



Transmon Implementations

Martin Leib, P. Zoller, WL, arXiv:1604.02359(2016).




- rotating frame
- only **pair interactions**
- **no crossings**
- all **local fields** have the **same sign**
- all **interactions** have the **same sign**
- programming [0,2]

$$\sigma_z^{(1)} \sigma_z^{(2)} \sigma_z^{(3)} \rightarrow \sigma_z^{(1)} \sigma_z^{(2)} + \sigma_z^{(2)} \sigma_z^{(3)} + \sigma_z^{(3)} \sigma_z^{(1)} + \sum_{i=1}^3 [2\sigma_z^{(i)} \sigma_z^{(a)} - \sigma_z^{(i)}] - 2\sigma_z^{(a)}$$


Quantum Circuit for LHZ

e.g. Flux-Qubit




Nicholas Chancellor, Stefan Zohren, Paul A. Warburton
arXiv:1603.09521 (2016).

e.g. Driven fixed frequency Transmons



D. C. McKay, S. Filipp, A. Mezzacapo, F. Solgun,
J. Chow, and J. M. Gambetta. arXiv:1604.03076 (2016).

e.g. Driven Kerr-nonlinearities



Shruti Puri, Christian Kraglund Andersen, Arne L. Grimsmo, Alexandre Blais, arXiv:1609.07117 (2016).